



Quiz #1: Immunization

True **False**

- 1. Mild illness is a reason to withhold vaccination.
- 2. If a mother is breastfeeding, she shouldn't be vaccinated.
- 3. A child's temperature should be checked routinely before vaccinations are administered.
- 4. A pregnancy test should be performed routinely for adolescent females before giving them MMR or varicella vaccines.
- 5. If there is an immunosuppressed child in the household, siblings should be given MMR and varicella vaccines.
- 6. If the first dose of hepatitis B vaccine was given more than one year ago, and no subsequent doses were given, you should repeat the first dose.
- 7. MMR and varicella vaccines can be given to a child whose mother is pregnant.
- 8. When a teen or adult has an injury and needs protection against tetanus, Td or Tdap is the correct vaccine choice, not Tt (tetanus toxoid) alone.
- 9. All health careworkers who have contact with patients and who have no contraindications should receive influenza vaccine every year.

Test Answers

1-F 2-F 3-F 4-F 5-T 6-F 7-T 8-T 9-T

If you missed any of these questions, you can find explanations for the answers in CDC's *Epidemiology and Prevention of Vaccine-Preventable Diseases* (a.k.a. "The Pink Book"), which you can download from CDC's website at www.cdc.gov/nip/publications/pink. You can also find information in the American Academy of Pediatrics current *Red Book*, which you can order online at www.aap.org/bookstore or by calling (847) 434-4000.

www.immunize.org/catg.d/p7001qz.pdf • Item # P7001 (4/06)